

This Page Is Inserted by IFW Operations
and is not a part of the Official Record

BEST AVAILABLE IMAGES

Defective images within this document are accurate representations of the original documents submitted by the applicant.

Defects in the images may include (but are not limited to):

- BLACK BORDERS
- TEXT CUT OFF AT TOP, BOTTOM OR SIDES
- FADED TEXT
- ILLEGIBLE TEXT
- SKEWED/SLANTED IMAGES
- COLORED PHOTOS
- BLACK OR VERY BLACK AND WHITE DARK PHOTOS
- GRAY SCALE DOCUMENTS

IMAGES ARE BEST AVAILABLE COPY.

**As rescanning documents *will not* correct images,
please do not report the images to the
Image Problem Mailbox.**

THIS PAGE BLANK (USPTO)

(57) Abstract: A hard surface antimicrobial cleaner is disclosed. The hard surface antimicrobial cleaner includes a disinfectant and a polysiloxane with at least one poly(oxyalkylene) side chain wherein the cleaner kills bacteria on a hard surface for a least 24 hours after being sprayed onto and wiped from the hard surface. The surface antimicrobial cleaner may include a solvent, a sequesterant, a surfactant, or a water soluble organosilane. Another version of the cleaner also includes a disinfectant and a polysiloxane with at least one poly(oxyalkylene) side chain and inhibits biofilm formation on a hard surface for at least 24 hours after being sprayed onto the hard surface.



— before the expiration of the time limit for amending the claims and to be republished in the event of receipt of amendments

For two-letter codes and other abbreviations, refer to the "Guidance Notes on Codes and Abbreviations" appearing at the beginning of each regular issue of the PCT Gazette.

Hard Surface Antimicrobial Cleaner With Residual Antimicrobial Effect

CROSS-REFERENCES TO RELATED APPLICATIONS

[0001] This application claims the benefit of United States Provisional Patent Application No. 60/275,405 filed March 13, 2001.

STATEMENT REGARDING FEDERALLY SPONSORED RESEARCH

5 **[0002]** Not Applicable.

BACKGROUND OF THE INVENTION

1. Field of the Invention

10 **[0003]** This invention relates to a hard surface antimicrobial cleaner with a residual antimicrobial effect, and to hard surface antimicrobial cleaner that inhibits the formation of biofilm on the hard surface.

2. Description of the Related Art

15 **[0004]** It is well known that the washing of hard surfaces (e.g., glass, tile, porcelain, fiberglass composites, metallic surfaces, ceramic surfaces, laminate surfaces, hard polymeric surfaces) with antimicrobial cleaners can remove many bacteria from the washed surfaces. Removal of the bacteria may be due to surfactants or disinfectants in the cleaner and/or the mechanical action of the wash procedure. Antimicrobial hard surface cleaners have been marketed in a variety of forms for some time. Typically, these hard surface antimicrobial products have been formulated to provide bacteria removal during washing.

20 However, there has been more interest recently in hard surface cleaners that have also been shown to provide a residual effectiveness against bacteria. By residual effectiveness it is meant that bacteria on a surface is killed for some period of time following the washing process. Given the potential severe health impacts of bacteria, there is a continuing search for improved antimicrobial cleaners which provide residual effectiveness versus bacteria.

25 **[0005]** Hard surfaces may also be prone to the attachment of biofilm, which also may have health impacts. A biofilm consists of cells immobilized on a surface and embedded in an organic polymer matrix of microbial origin. A biofilm

is a surface accumulation, which is not necessarily uniform in time or space. A biofilm may be composed of a significant fraction of inorganic or abiotic substances held cohesively by the biotic matrix. A biofilm is a protective matrix for bacteria, with the essential purpose of survival in an environment of limited nutrient supply. Biofilms consist of both host microbes and their extracellular products, usually exopolysaccharides. Microbes have a tendency to form these protective exopolysaccharide matrices after they have adhered to a surface. The formation of biofilm complexes requires only humid conditions and/or water systems and contact with a support surface. With respect to nutrients, a nutrient deficiency in fact may increase the biofilm formation capacity of microbes.

[0006] Biofilms generally can be produced by almost all microbes under suitable conditions. The most common biofilm producers belong to the genera *Pseudomonas*, *Enterobacter*, *Flavobacterium*, *Alcaligenes*, *Staphylococcus*, *Klebsiella* and *Bacillus*. One of the main purposes of natural biofilm formation is for the protection of the host microbes from a hostile environment. As a consequence, there is a combative interaction between microbes in biofilms and biocides such as disinfectants. Further, the sessile mode of bacterial growth in biofilms differs from that of the same bacteria species that are present as planktonic cells in a circulating aqueous medium which interfaces with the biofilm. Because of the ramifications of biofilm formation, there have been proposed techniques to inhibit the growth of biofilm on a surface. For example, surfactants have been added to aqueous systems to inhibit microbial colonization on a surface (see, e.g., U.S. Patent No. 6,039,965). However, there is a need for a hard surface antimicrobial cleaner that cleans and disinfects a hard surface and thereafter inhibits the formation of biofilm on the hard surface.

[0007] Thus, given the potential health impacts of bacteria and biofilm on a surface, particularly in kitchen areas, there is a need for hard surface antimicrobial cleaners which clean and disinfect a hard surface and also provide for residual effectiveness versus bacteria. Further, there is a need for hard surface antimicrobial cleaners which clean and disinfect a hard surface and thereafter inhibit the formation of biofilm on the hard surface.

BRIEF SUMMARY OF THE INVENTION

[0008] The foregoing needs are met by a hard surface antimicrobial cleaner according to the invention including a disinfectant and a polysiloxane with at least one poly(oxyalkylene) side chain wherein the cleaner kills bacteria on a hard surface for at least 24 hours after being sprayed onto and wiped from the hard surface. The polysiloxane has the formula:



in which R_1 , R_2 , R_4 , R_5 , R_6 are identical or different and are a C_1 - C_6 alkyl or phenyl, R_3 is identical or different and is C_1 - C_6 alkyl, phenyl or Q, and Q is an ether polyoxyalkylene group of the formula $-R-O-(R'O)_nR''$ where R is a linear or branched C_3 - C_{15} alkyl group, $(R'O)_n$ is a poly(ethyleneoxy) and/or poly(propyleneoxy) group, n is a mean value ranging from 5 to 200, R'' is H or a C_1 - C_6 alkyl group, p is a mean value ranging from 10 to 200, and q is 0 or a mean value ranging from 1 to 200, R_3 being Q when q is 0.

[0009] In a first version of the invention, the hard surface antimicrobial cleaner includes from about 0.01% to about 20% by weight of the total weight of the cleaner of the disinfectant; from about 0.01% to about 20% by weight of the total weight of the cleaner of the polysiloxane with at least one poly(oxyalkylene) side chain; and from about 0.5% to about 20% by weight of the total weight of the cleaner of a solvent.

[0010] In a second version of the invention, the hard surface antimicrobial cleaner includes from about 0.01% to about 20% by weight of the total weight of the cleaner of the disinfectant; from about 0.01% to about 20% by weight of the total weight of the cleaner of the polysiloxane with at least one poly(oxyalkylene) side chain; and from 0.1 to about 20% by weight of the total weight of the cleaner of a sequesterant.

[0011] In a third version of the invention, the hard surface antimicrobial cleaner includes from about 0.01% to about 20% by weight of the total weight of the cleaner of the disinfectant; from about 0.01% to about 20% by weight of the total weight of the cleaner of the polysiloxane with at least one poly(oxyalkylene) side chain; and from 0.5 to about 20% by weight of the total weight of the cleaner of a surfactant selected from nonionic surfactants, amphoteric surfactants, sarcosine anionic surfactants, cationic surfactants and mixtures thereof.

[0012] In a fourth version of the invention, the hard surface antimicrobial cleaner includes from about 0.01% to about 20% by weight of the total weight of the cleaner of the disinfectant; from about 0.01% to about 20% by weight of the total weight of the cleaner of the polysiloxane with at least one poly(oxyalkylene) side chain; and an organosilane of the formula: $A_{3-x}B_xSiD$ wherein A is -OH or a hydrolyzable group, B is an alkyl group of from 1 to 4 carbon atoms, x has a value of 0, 1 or 2, and D is a hydrocarbon group of from 1 to 4 carbon atoms, phenyl, or a nonionic or cationic, substituted-hydrocarbon group containing at least one oxygen or nitrogen group or salts of such substituted-hydrocarbon groups.

[0013] In a fifth version of the invention, the hard surface antimicrobial cleaner includes from about 50% to about 99.9% by weight of the total weight of the cleaner of an alkyl alcohol disinfectant; and the polysiloxane with at least one poly(oxyalkylene) side chain, wherein the cleaner inhibits biofilm formation on a hard surface for at least 24 hours after being sprayed onto the hard surface. In the fifth version of the invention, the alkyl alcohol disinfectant is preferably ethanol, n-propanol or isopropanol, and serves to disinfect the hard surface and to distribute the polysiloxane on the hard surface in order to inhibit biofilm formation on the hard surface. All or most of the alkyl alcohol disinfectant may eventually evaporate from the hard surface due to the volatility of the alcohol. The fifth version of the invention may consist essentially of the alkyl alcohol disinfectant, the polysiloxane and water, if desired. Alternatively, the fifth version of the invention may include a solvent, a sequesterant, a surfactant or an organosilane.

[0014] It is therefore an advantage of the present invention to provide a hard surface antimicrobial cleaner that may be used to clean and disinfect a hard surface and also provides for residual effectiveness versus bacteria.

[0015] It is another advantage of the present invention to provide a hard surface antimicrobial cleaner that may be used to clean and disinfect a hard surface and thereafter inhibits the formation of biofilm on the hard surface.

[0016] These and other features, aspects, and advantages of the present invention will become better understood upon consideration of the following detailed description and appended claims.

DETAILED DESCRIPTION

5 [0017] An antimicrobial cleaner according to the invention will be useful for all hard surface cleaning and disinfectant formulations, including kitchen cleaners and disinfectants, bathroom cleaners and disinfectants, all-purpose cleaners and disinfectants, toilet cleaners and disinfectants (both periodic and continuous), bowl
10 cleaners and disinfectants, and drain cleaners and disinfectants. An antimicrobial cleaner according to the invention is particularly suitable as a fully diluted hard surface cleaner. As such, it can be used without further dilution by applying it at full strength to a soiled hard surface, and wiping and/or scrubbing to remove the
10 soil.

[0018] The cleaner is especially useful for cleaning kitchen surfaces which are soiled. For instance, when certain embodiments of the invention are used as a kitchen cleaner and disinfectant, the cleaner would have the advantage of killing
15 bacteria for at least 24 hours after application (i.e., the cleaner provides a residual effectiveness against bacteria). Of course, upon spillage of containment such as gravy on a cleaned kitchen surface, one would re-use the cleaner.

[0019] When certain other embodiments of the invention are used as a hard surface cleaner and disinfectant, the cleaner would have the advantage of inhibiting the formation of biofilm on the hard surface for at least 24 hours after
20 application. In other words, the cleaner provides a residual inhibition of the formation of biofilm. By residual inhibition, it is meant that biofilm does not form on the surface for some period of time following the application process.




[0020] A hard surface antimicrobial cleaner according to the invention includes from about 0.01% to about 99% by weight of the total weight of the cleaner of a
25 disinfectant (as 100% active). Suitable disinfectants include, for example, quaternary ammonium compounds, phenolics (aromatic alcohols), guanide derivatives, ampholytes (betaines), aldehydes (such as glutaraldehyde and formaldehyde), and alkyl alcohols. A disinfectant can be understood to be a hygiene agent which shows a reduction in the number of viable microorganisms in
30 a specified culture when used at a specified level. In one embodiment of the hard surface antimicrobial cleaner, the disinfectant is other than an alkyl alcohol, and the cleaner preferably includes from about 0.01% to about 20% by weight of the total weight of the cleaner of the disinfectant, and most preferably from about

0.1% to about 2% by weight of the total weight of the cleaner of the disinfectant. In another embodiment of the hard surface antimicrobial cleaner, the disinfectant is a volatile alkyl alcohol such as ethanol or propanol, and the cleaner preferably includes from about 50% to about 99.9% by weight of the total weight of the cleaner of the alkyl alcohol, and most preferably from about 60% to about 80% by weight of the total weight of the cleaner of the alkyl alcohol.

[0021] Non-limiting illustrative disinfectant quaternary ammonium compounds include benzalkonium chloride, alkyl-dimethyl-benzylammonium chloride, alkyl-dimethyl-ethylbenzylammonium chloride, dodecyl-dimethyl-3,4-dichlorobenzylammonium chloride, dodecyl-di-(2-hydroxyethyl)-benzylammonium chloride, 4-diisobutyl-phenoxyethoxyethyl-dimethylbenzylammonium chloride, 4-diisobutyl-cresoxyethoxyethyl-dimethylbenzylammonium chloride, dimethyl-didecylammonium chloride, cetyl-trimethylammonium bromide, dodecyl-pyridinium chloride, cetyl pyridinium chloride, dodecyl-isoquinolinium chloride, decamethylene-bis-4-aminoquinaldinium dichloride, and mixtures thereof. One example quaternary ammonium compound is BTC 2125M, an alkyldimethylbenzyl ammonium chloride and dimethyl ethylbenzyl ammonium chloride mixture commercially available from Stepan.

[0022] Non-limiting illustrative disinfectant phenolics include phenol, mono- and poly-chlorophenols, cresols, 4-chloro-3-methylphenol, 3,5-dimethyl-4-chlorophenol, thymol, 4-chlorothymol, 4-t-amylphenol, saligenin, 4-n-hexylresorcinol, carvacrol, 2-phenylphenol, 2-benzyl-4-chlorophenol, 2,2'-dihydroxy-5,5'-dichlorodiphenylmethane, 2,2'-dihydroxy-3,3',5,5',6,6'-hexachlorodiphenylmethane, 2,2'-dihydroxy-5,5'-dichlorodiphenyl sulphide, 2,2'-dihydroxy-3,3',5,5'-tetrachlorodiphenyl sulphide, 2-hydroxy-2',4,4'-trichlorodiphenyl ether, dibromosalicyl and mixtures thereof.

[0023] Non-limiting illustrative disinfectant guanide compounds include compounds of the general formula (I), (II) or (III) given below:



In the formulas (I), (II) and (III), X is an alkyl group, an aminoalkyl group, a phenyl group, an alkylphenyl group, a halophenyl group, a hydroxyphenyl group, a methoxyphenyl group, a carboxyphenyl group, a naphthyl group or a nitrile group; X' is a hydrogen atom or an alkyl group; and j and k each is a positive integer, preferably an integer within the range of 2 to 10. A preferred example of a suitable guanide compound is chlorhexidine, also known as 1,6-bis(N⁵-p-chlorophenyl-N¹-biguanido)hexane.


[0024] Non-limiting illustrative disinfectant alkyl alcohols include ethanol, n-propanol, isopropanol and mixtures thereof.

[0025] A hard surface antimicrobial cleaner according to the invention includes from about 0.01% to about 20% by weight of the total weight of the cleaner of a polysiloxane with at least one poly(oxyalkylene) side chain. Preferably, the polysiloxane has the formula: $\text{R}_1\text{R}_2\text{R}_3\text{SiO}(\text{R}_4\text{R}_5\text{SiO})_p(\text{R}_6\text{QSiO})_q\text{SiR}_3\text{R}_2\text{R}_1$ in which R₁, R₂, R₄, R₅, R₆ are identical or different and are a C₁-C₆ alkyl or phenyl, R₃ is identical or different and is C₁-C₆ alkyl, phenyl or Q, and Q is an ether polyoxyalkylene group of the formula $-\text{R}-\text{O}-(\text{R}'\text{O})_n\text{R}''$ where R is a linear or branched C₃-C₁₅ alkyl group, (R'O)_n is a poly(ethyleneoxy) and/or poly(propyleneoxy) group, n is a mean value ranging from 5 to 200, R'' is H or a C₁-C₆ alkyl group, p is a mean value ranging from 10 to 200, and q is 0 or a mean

value ranging from 1 to 200, R_3 being Q when q is 0. These polymers are shown in PCT International Publication WO 99/18784 which is incorporated herein by reference. One preferred polysiloxane will be referred to as "S2" and has the formula:

5

10



15

20

Polysiloxanes with one or more poly(oxyalkylene) side chains can be made by processes described in U.S. Patent Nos. 6,337,383 and 3,172,899 which are incorporated herein by reference. A hard surface antimicrobial cleaner according to the invention preferably includes from about 0.01% to about 5% by weight of the total weight of the cleaner of the polysiloxane. A hard surface antimicrobial cleaner according to the invention most preferably includes from about 0.025% to about 1% by weight of the total weight of the cleaner of the polysiloxane (as 100% active).

25

30

[0026] A hard surface antimicrobial cleaner according to the invention may optionally include from 0 to about 20% by weight of the total weight of the cleaner of a surfactant (as 100% active) which can be a nonionic surfactant, an amphoteric surfactant, a sarcosine anionic surfactant, a cationic surfactant or mixtures thereof. Preferably, the surfactant is used in an amount of 0.5% to 10% by weight of the total weight of the cleaner. More preferably, the surfactant is used in an amount of 1% to 2% by weight of the total weight of the cleaner. The term "amphoteric" surfactant includes "zwitterionic" surfactants for the purposes of this invention since those terms are often used almost interchangeably. These surfactants are well known and a large number are commercially available as can be seen from an examination of the widely available "McCutcheon's Emulsifiers & Detergents" and the "CTFA Cosmetic Ingredient Dictionary".

35

[0027] Examples of nonionic surfactants include alcohol ethoxylates such as C_8 to C_{18} alcohol ethoxylates containing from about 3 to 50 moles of ethylene oxide per molecule; C_8 to C_{18} fatty acid esters and amides containing from about 2

to 50 moles of ethylene oxide; C₈ to C₁₈ fatty alcohols; C₈ to C₁₈ diols such as tetramethyl decynediol and dimethyl octynediol; block copolymers of polyethylene oxide and polypropylene oxide; C₈ to C₁₈ fatty acid esters of glycerine; ethoxylated and propoxylated C₈ to C₁₈ fatty alcohols; C₈ to C₁₈ fatty amine and amidoamine oxides; C₈ to C₁₈ fatty amides and alkanolamides; and alkyl saccharides (e.g., alkyl glucosides) or alkenyl-saccharides such as a saccharide having the formula: R₁₀-O-(R₁₂O)_t-(G)_p where R₁₀ is a linear or branched alkyl, alkenyl or alkyl-phenyl group having 6-18 carbon atoms, R₁₂ is an alkylene group having 2-4 carbon atoms, G is a reduced saccharide residue having 5-6 carbon atoms, t is 0-10, and p is 1-10.

[0028] Examples of amphoteric surfactants include amine oxides, C₈ to C₁₈ sultaines such as coco-sultaine and cocamidopropyl hydroxysultaine; C₈ to C₁₈ fatty derivatives of amino acids such as cocamphocarboxyglycinate and lauramphoglycinate; C₈ to C₁₈ alkyl betaines such as decyl betaine, coco-betaine, lauryl betaine, myristyl betaine and stearyl betaine; and C₈ to C₁₈ amidoalkyl betaines such as cocoamidoethyl betaine, cocamidopropyl betaine, lauramidopropyl betaine, myristamidopropyl betaine and oleamidopropyl betaine.

[0029] Since sarcosine surfactants are known to be compatible with quaternary ammonium compounds, this class of anionic surfactants can be used with quaternary ammonium compounds. Examples of such surfactants are C₈ to C₁₈ alkyl sarcosines and their alkali metal or ammonium salts such as sodium, potassium, lithium or ammonium C₈ to C₁₈ alkyl sarcosinates.

[0030] Examples of cationic surfactants other than the quaternary ammonium compounds already described above are quaternary ammonium compounds which contain at least two nitrogen-bonded alkyl chains having at least about 16 carbon atoms such as distearyldimonium chloride and ditallowdimonium chloride; C₈ to C₁₈ fatty alkyl amines, amidoalkylamines and amidoalkanolamines, and their salts; ethoxylated amines; amine oxides; and imidazoline.

[0031] A hard surface antimicrobial cleaner according to the invention may optionally include from 0 to about 20% by weight of the total weight of the cleaner of a sequesterant (as 100% active) such as ethylenediamine tetraacetic acid (EDTA) or its salts (e.g. EDTA, sodium salt), phosphonates, nitrilotriacetic acid (NTA) or its salts, hydroxyethylene diamine and triacetic acid (HEDTA) or its salts,

and diethylene triamine pentaacetic acid (DTPA) or its salts. Preferably, the sequesterant is used in an amount of 0.1% to 15% by weight of the total weight of the cleaner. More preferably, the sequesterant is used in an amount of 0.05% to 5% by weight of the total weight of the cleaner.

5 **[0032]** A hard surface antimicrobial cleaner according to the invention may optionally include from 0 to about 20% by weight of the total weight of the cleaner of a solvent (as 100% active) such as alcohols, glycols, ethers and glycol ethers, such as diethylene glycol monobutyl ether ("Butyl Carbitol"), dipropylene glycol n-butyl ether ("DPnB"), propylene glycol n butyl ether ("PnB"), ethylene glycol butyl
10 ether ("Butyl Cellosolve"), dipropylene glycol monomethylether, propylene glycol, carbitol, methoxypropanol, glycerine, isopropanol and ethanol. Preferably, the solvent is used in an amount of 0.5% to 10% by weight of the total weight of the cleaner. Most preferably, the solvent is used in an amount of 0.5% to 3% by weight of the total weight of the cleaner.

15 **[0033]** A hard surface antimicrobial cleaner according to the invention may optionally include from 0 to about 5% by weight of the total weight of the cleaner of an organosilane of the formula: $A_{3-x}B_xSiD$ wherein A is -OH or a hydrolyzable group, B is an alkyl group of from 1 to 4 carbon atoms, x has a value of 0, 1 or 2, and D is a hydrocarbon group of from 1 to 4 carbon atoms, phenyl, or a nonionic
20 or cationic, substituted-hydrocarbon group containing at least one oxygen or nitrogen group or salts of such substituted-hydrocarbon groups. These organosilanes and methods for stabilizing these organosilanes are described in U.S. Patent Nos. 6,087,319 and 5,411,585 which are incorporated herein by reference. Preferably, A is selected from the group consisting of -OR¹ and -OR^{2A}
25 OR¹ where each R¹ is R² or hydrogen, R² is an alkyl group of 1 to 4 carbon atoms, R^{2A} is a divalent saturated hydrocarbon group of from 1 to 4 carbon atoms, x has a value of 0 or 1, and D is selected from the group consisting of alkyl groups of from 1 to 4 carbon atoms, vinyl, allyl, glycidoxypentyl, -R³N(R⁴)_y H_{2-y}, -R³N⁽⁺⁾(R⁴)_y H_{3-y} X⁽⁻⁾, -R³NHR³N(R⁴)_yH_{2-y}, -R³NHR³N⁽⁺⁾(R⁴)_yH_{3-y} X⁽⁻⁾, -R³N⁽⁺⁾R²R⁴R⁵X⁽⁻⁾ wherein
30 R³ is a divalent saturated hydrocarbon group of from 1 to 12 carbon atoms; R⁴ and R⁵ are each selected from the group consisting of alkyl groups of 1 to 18 carbon atoms, -CH₂C₆H₅, -CH₂CH₂OH and -CH₂OH; y has a value of 0, 1 or 2; and X is an anion. Most preferably, the organosilane is

3-trimethoxysilyl)propyldimethyloctadecyl ammonium chloride (commercially available as Dow Corning 5772) or

3-trimethoxysilyl)propylmethyldi(decyl)ammonium chloride. Preferably, a hard surface antimicrobial cleaner according to the invention includes from 0.01 to about 3% by weight of the total weight of the cleaner of any of the above organosilanes. The organosilane can further enhance the residual effectiveness against bacteria.

[0034] Optionally, other additives such as pH adjusters, buffers, detergent builders, acids, dyes, fragrance, viscosity adjusters, and corrosion inhibitors can be included in the hard surface antimicrobial cleaner of the present invention provided that they are compatible with the other ingredients. The hard surface antimicrobial cleaner of the present invention is typically formulated as an aqueous solution; however, water is not required in the cleaner. For example, the cleaner can be supplied as a concentrate, or water can be left out of the cleaner in favor of an alternative solvent such as an alkyl alcohol. A non-concentrated formula would be as described above.

Examples

[0035] The following examples serve to further illustrate the invention. The examples are not intended to limit the invention in any way.

Example 1

[0036] A formulation using the ingredients of Table 1 was prepared.

Table 1

Chemical Description	Wt. %	Functional Description	Predicted Function within the formulation
Deionized Water	96.350		
EDTA tetra sodium salt (40%)	0.400	sequesterant	boosts cleaning by removing calcium soils and deposits
Ninox DO40 C ₁₀ dimethyl amine oxide 40%	0.563	surfactant	emulsifies soils
Tergitol 15-S-7 (alcohol ethoxylate)	0.300	surfactant	emulsifies soils
Glucopon 425N (Alkyl Glucoside)	0.563	surfactant	emulsifies soils
Catigene 4513-80% Europe or BTC 2125 M (80%) US quaternary ammonium disinfectant	0.413	disinfectant	kills microbes
Butyl Cellosolve (ethylene glycol monobutyl ether)	0.751	solvent	removes soiling
Polymer S2	0.050	disinfectant enhancer	retains disinfectant on treated surfaces
Sodium Hydroxide (30%)	0.360	alkali	pH balance
Fragrance IFF 4640 HBD (Lemon)	0.250	fragrance	fragrance

Example 2

[0037] A formulation using the ingredients of Table 2 was prepared.

Table 2

Chemical Description	Wt. %	Functional Description	Predicted Function within the formulation
Deionized Water	96.450		
EDTA tetra sodium salt (40%)	0.400	sequesterant	boosts cleaning by removing calcium soils and deposits
Ninox DO40 C ₁₀ dimethyl amine oxide 40%	0.563	surfactant	emulsifies soils
Tergitol 15-S-7 (alcohol ethoxylate)	0.300	surfactant	emulsifies soils
Glucopon 425N (Alkyl Glucoside)	0.563	surfactant	emulsifies soils
Catigene 4513-80% Europe or BTC 2125 M 80% US quaternary ammonium disinfectant	0.413	disinfectant	kills microbes
Butyl Cellosolve (ethylene glycol monobutyl ether)	0.751	solvent	Removes soiling
Polymer S2	0.050	disinfectant enhancer	retains disinfectant on treated surfaces
Sodium Hydroxide (30%)	0.360	alkali	pH balance
Fragrance Takasago RM-1489	0.150	fragrance	fragrance

Example 3

[0038] A formulation using the ingredients of Table 3 was prepared.

Table 3

Chemical Description	Wt. %	Functional Description	Predicted Function within the formulation
Deionized Water	95.437		
EDTA tetra sodium salt (40%)	0.400	sequesterant	boosts cleaning by removing calcium soils and deposits
Ninox DO 40 (C10 dimethyl amine oxide) (40%)	0.750	surfactant	emulsifies soils
Tergitol 15-S-7 (alcohol ethoxylate)	0.200	surfactant	emulsifies soils
Glucopon 425N (Alkyl Glucoside)	0.750	surfactant	emulsifies soils
Catigene 4513-80% Europe or BTC 2125 M (80%) USA (quaternary ammonium disinfectant)	0.550	disinfectant	Kills microbes
Butyl Cellosolve (ethylene glycol monobutyl ether)	0.570	solvent	removes soiling
Hexyl Cellosolve (ethylene glycol n-hexyl ether)	0.430	solvent	removes soiling
Polymer S2	0.300	disinfectant enhancer	retains disinfectant on treated surfaces
Sodium Hydroxide (30%)	0.363	Alkali	pH balance
Fragrance IFF 4640 HBD (Lemon)	0.250	fragrance	fragrance

Example 4

[0039] The formulations of Examples 1 and 3 were tested against a competitive product with a claim of residual antimicrobial activity. To measure the residual antimicrobial benefit of the formulations of Examples 1 and 3, surfaces were treated with the formulation according to label directions. After 24 hours, under normal room conditions, surfaces were wiped and inoculated with appropriate test organisms. Residual activity was measured by log reduction of test organism compared to parallel controls.

[0040] The following results were obtained as shown in Table 4. Both Example 1 and Example 3 formulas were superior to the competitor product.

Table 4

Residual Antibacterial Results		
Test Organism is <i>Staphylococcus aureus</i> (Staph. Aureus)		
Test Surface is Glazed Ceramic Tile		
Log reductions based on a comparison with log recovery of Controls.		
Controls are Glazed Ceramic Tiles treated with 0.01% Triton X-100 surfactant		
Formula	Bacterial Log Reduction	Bacterial Log Reduction
Example 1	4.17	
Example 3		3.51
Competitor Product with Residual Claim	0.13	0.39
Controls (Log recovery)	6.64	6.93

Example 5

[0041] A formulation using the ingredients of Table 5 was prepared.

5

Table 5

Chemical Description	Wt. %	Functional Description	Predicted Function within the formulation
Deionized Water	96.305		
EDTA tetra sodium salt (40%)	0.400	sequesterant	boosts cleaning by removing calcium soils and deposits
Ammonyx DO 40 C10 dimethyl amine oxide	0.750	surfactant	emulsifies soils
Tergitol 15-S-7 (alcohol ethoxylate)	0.300	surfactant	emulsifies soils
Glucopon 425N (Alkyl Glucoside)	0.563	surfactant	emulsifies soils
BTC 2125 M (quaternary ammonium disinfectant)	0.413	disinfectant	kills microbes
Butyl Cellosolve (ethylene glycol monobutyl ether)	0.751	solvent	removes soiling
Polymer S2	0.050	disinfectant enhancer	retains disinfectant on treated surfaces
Sodium Hydroxide (30%)	0.218	Alkali	pH balance
Fragrance IFF 4641 HBD	0.250	fragrance	Fragrance

Example 5a

[0042] Using gravimetric analysis, the formulation of Example 5 was analyzed by measuring the percent removal of 25-29 milligrams of a lab generated, synthetic, greasy kitchen soil from enamel coated steel tiles with a Gardner Scrub Machine after 7 strokes. The percent removal was 94.4% soil removal.

10

Example 5b

[0043] The formulation of Example 5 was tested to measure the residual antimicrobial benefit of the formulation on surfaces treated with the formulation. After a test time, under normal room conditions, surfaces were wiped and inoculated with appropriate test organisms. Residual activity was measured by log reduction of test organism. The results are in Table 5b below:

Table 5b

Test No.	Test Surface	Test Condition	Organism	Test time	Log Reduction
1	ceramic tile	Dry Wipe	<i>Staph. Aureus</i>	24 hours	3.30
2	ceramic tile	Dry Wipe	<i>Staph. Aureus</i>	24 hours	3.72
3	ceramic tile	Dry Wipe	<i>Staph. Aureus</i>	12 Hours	4.81 *
4	ceramic tile	Dry Wipe	<i>Staph. Aureus</i>	12 hours	5.87 **
5	Stainless steel	Dry Wipe	<i>Staph. Aureus</i>	24 hours	1.50
6	Formica	Dry Wipe	<i>Staph. Aureus</i>	24 hours	4.55
7	Stainless steel	Dry Wipe	<i>Staph. Aureus</i>	24 hours	> 5.74
8	Formica	Dry Wipe	<i>Staph. Aureus</i>	24 hours	> 5.75
* with an organic soil load (0.03% bovine serum) ** without an organic soil load					

Example 6

[0044] A formulation using the ingredients of Table 6 was prepared.

Table 6

Chemical Description	Wt. %	Functional Description	Predicted Function within the formulation
Deionized Water	96.130		
EDTA tetra sodium salt (40%)	0.400	sequesterant	boosts cleaning by removing calcium soils and deposits
Ammonyx DO 40 C10 dimethyl amine oxide	0.750	surfactant	emulsifies soils
Tergitol 15-S-7 (alcohol ethoxylate)	0.300	surfactant	emulsifies soils
Glucopon 425N (Alkyl Glucoside)	0.563	surfactant	emulsifies soils
BTC 2125 M (quaternary ammonium disinfectant)	0.413	disinfectant	kills microbes
Butyl Cellosolve (ethylene glycol monobutyl ether)	0.751	Solvent	removes soiling
Polymer S2	0.225	disinfectant enhancer	retains disinfectant on treated surfaces
Sodium Hydroxide (30%)	0.218	Alkali	pH balance
Fragrance IFF 4641 HBD	0.250	fragrance	Fragrance

Example 6a

[0045] Using gravimetric analysis, the formulation of Example 6 was analyzed by measuring the percent removal of 25-29 milligrams of a lab generated, synthetic, greasy kitchen soil from enamel coated steel tiles with a Gardner Scrub Machine after 7 strokes. The percent removal was 92% soil removal.

Example 6b

[0046] The formulation of Example 6 was tested to measure the residual antimicrobial benefit of the formulation on surfaces treated with the formulation. After a test time, under normal room conditions, surfaces were wiped and inoculated with appropriate test organisms. Residual activity was measured by log reduction of test organism. The results are in Table 6b below:

Table 6b

Test No.	Test Surface	Test Condition	Organism	Test time	Log Reduction
1	Glass slides	Dry Wipe	<i>Staph. Aureus</i>	24 hours	3.52

Example 7

[0047] A formulation using the ingredients of Table 7 was prepared.

Table 7

Chemical Description	Wt. %	Functional Description	Predicted Function within the formulation
Deionized Water	96.255		
EDTA tetra sodium salt (40%)	0.400	sequesterant	boosts cleaning by removing calcium soils and deposits
Ammonyx DO 40 C10 dimethyl amine oxide	0.750	surfactant	emulsifies soils
Tergitol 15-S-7 (alcohol ethoxylate)	0.300	surfactant	emulsifies soils
Glucopon 425N (Alkyl Glucoside)	0.563	surfactant	emulsifies soils
BTC 2125 M (quaternary ammonium disinfectant)	0.413	disinfectant	Kills microbes
Butyl Cellosolve (ethylene glycol monobutyl ether)	0.751	solvent	removes soiling
Polymer S2	0.050	disinfectant enhancer	retains disinfectant on treated surfaces
3-trimethoxysilyl)propyldimethyloctadecyl ammonium chloride	0.050	quaternary organosilane	Extra residual biocidal performance
Sodium Hydroxide (30%)	0.218	Alkali	pH balance
Fragrance IFF 4641 HBD	0.250	fragrance	Fragrance

Example 7a

[0048] Using gravimetric analysis, the formulation of Example 7 was analyzed by measuring the percent removal of 25-29 milligrams of a lab generated, synthetic, greasy kitchen soil from enamel coated steel tiles with a Gardner Scrub Machine after 7 strokes. The percent removal was 91.6% soil removal.

Example 7b

[0049] The formulation of Example 7 was tested to measure the residual antimicrobial benefit of the formulation on surfaces treated with the formulation. After a test time, under normal room conditions, surfaces were wiped and inoculated with appropriate test organisms. Residual activity was measured by log reduction of test organism. The results are in Table 7b below:

Table 7b

Test No.	Test Surface	Test Condition	Organism	Test time	Log Reduction
1	Ceramic tile	Dry Wipe	<i>Staph. Aureus</i>	24 hours	2.83
2	Ceramic tile	Dry Wipe	<i>Staph. Aureus</i>	24 hours	4.87

Example 8

[0050] A formulation using the ingredients of Table 8 was prepared.

Table 8

Chemical Description	Wt. %	Functional Description	Predicted Function within the formulation
Deionized Water	76.1380		
EDTA tetra sodium salt (40%)	14.1620	sequesterant	boosts cleaning by removing calcium soils and deposits
Butyl Carbitol (diethylene glycol butyl ether)	5.8410	Solvent	removes soils
Tergitol NP-10	0.7302	surfactant	emulsifies soils
BTC 2125 M (quaternary ammonium disinfectant)	0.2633	disinfectant	kills microbes
EDTA acid	0.1809	Acid form of sequesterant	pH adjustment
Polymer S2	0.1460	disinfectant enhancer	retains disinfectant on treated surfaces
Rhodoquat RP50 (50%) (benzalkonium chloride)	2.4900	disinfectant	kills microbes
Givaudan-Roure PA 55386	0.0486	fragrance	Fragrance

Example 9

[0051] A formulation using the ingredients of Table 9 was prepared.

Table 9

Chemical Description	Wt. %	Functional Description	Predicted Function within the formulation
Deionized Water	73.25		
Butyl Carbitol (diethylene glycol butyl ether)	2.00	Solvent	removes soils
dipropylene glycol butyl ether	3.00	Solvent	removes soils
BTC 888 (quaternary ammonium disinfectant)	0.27	disinfectant	kills microbes
Plurafac B25-5 (alcohol ethoxylate)	1.25	surfactant	emulsifies soils
Ethomeen 0/12 (ethoxylated oleyl amine)	1.00	surfactant	emulsifies soils
Mackam 2CSF (disodium cocoamphodipropionate)	4.95	surfactant	emulsifies soils
EDTA tetra sodium salt (40%)	12.00	sequesterant	boosts cleaning by removing calcium soils and deposits
EDTA acid	0.13	Acid form of sequesterant	pH adjustment
Dequest 2016 (phosphonate)	1.40	Corrosion inhibitor	
AMP-95 (2-amino 2-methyl 1-propanol)	0.50	Solvent	Corrosion inhibitor
Polymer S2	0.20	disinfectant enhancer	retains disinfectant on treated surfaces
Givaudan-Roure PA 55386	0.05	fragrance	fragrance

5

Example 9a

[0052] Using gravimetric analysis, the formulation of Example 9 was analyzed by measuring the percent removal of a lab generated, soap scum from 6 ceramic tiles with a Gardner Scrub Machine after 7 strokes. The percent removal for 3 tests was 70%, 75.6% and 89.6% soil removal.

10

Example 9b

[0053] The formulation of Example 9 was tested to measure the residual antimicrobial benefit of the formulation on surfaces treated with the formulation. After a test time, under normal room conditions, surfaces were wiped and

15

inoculated with appropriate test organisms. Residual activity was measured by log reduction of test organism. The results are in Table 9b below:

Table 9b

Test No.	Test Surface	Test Condition	Organism	Test time	Log Reduction
1	Ceramic tile	Dry Wipe	<i>Staph. Aureus</i>	1 minute contact	2.63
2	Ceramic tile	Dry Wipe	<i>Staph. Aureus</i>	10 minute contact	4.28

5

Example 10

[0054] A formulation using the ingredients of Table 10 was prepared.

Table 10

Chemical Description	Wt. %	Functional Description	Predicted Function within the formulation
Deionized Water	73.15		
Butyl Carbitol (diethylene glycol butyl ether)	2.00	Solvent	removes soils
dipropylene glycol butyl ether	3.00	Solvent	removes soils
BTC 888 (quaternary ammonium disinfectant)	0.27	disinfectant	kills microbes
Plurafac B25-5 (alcohol ethoxylate)	1.25	surfactant	emulsifies soils
Ethomeen 0/12 (ethoxylated oleyl amine)	1.00	surfactant	emulsifies soils
Mackam 2CSF (disodium cocoamphodipropionate)	4.95	surfactant	emulsifies soils
EDTA tetra sodium salt (40%)	12.00	sequesterant	boosts cleaning by removing calcium soils and deposits
EDTA acid	0.13	Acid form of sequesterant	pH adjustment
Dequest 2016 (phosphonate)	1.40	Corrosion inhibitor	
AMP-95 (2-amino 2-methyl 1-propanol)	0.50	Solvent	Corrosion inhibitor
Polymer S2	0.20	disinfectant enhancer	retains disinfectant on treated surfaces
3-trimethoxysilyl)propyldimethyloctadecyl ammonium chloride	0.15	quaternary organosilane	Extra residual biocidal performance

10

Example 11

[0055] The Polymer S2 was made up in isopropanol at 2% w/v and diluted 2X serially to produce a 1% w/v solution, a 0.5% solution and a 0.25% w/v solution of the Polymer S2 in isopropanol. Two microliters of these materials were spread
5 over approximately 1 square inch of the surface of a sterile petri dish (polystyrene or glass). The polystyrene and glass surfaces were treated with the alcohol solution (0.25% w/v, 0.5% w/v, 1% w/v, 2% w/v) of the polymer S2 at a rate of 5 to 40 μ g/sq. in. respectively. The plates were dried at 35°C for 10-15 minutes to remove alcohol and leave a film of polymer. Fifteen milliliters of microbiological
10 growth medium (tryptic soy broth) were added to each plate. Each plate was placed on a reciprocating shaker overnight at 70 rpm at ambient temperature (18-20°C). After 18-20 hours, the medium was removed and replaced with fresh medium, and 100 microliters of a 24 hour culture of organisms *Klebsiella pneumonia* (gram negative bacteria) and *Pseudomonas fluorescens* (gram
15 negative bacteria) was inoculated into each plate. The plates were placed on the shaker overnight at 70 rpm. After 18-24 hours, the liquid culture was decanted, and the plates were washed with 3-15 ml. aliquots of tap water. The plates were dried and imaged as is or after staining with 10% Grams crystal violet for 1 minute, rinsed and dried. Macro and micro images were then taken to determine
20 the levels of biofilm formation in the (0.25% w/v, 0.5% w/v, 1% w/v, 2% w/v) plates. All plates showed virtually complete inhibition of *Klebsiella pneumonia* and *Pseudomonas fluorescens* biofilm formation.

[0056] The above procedure was also undertaken with a competitive product disinfectant aerosol spray having a claim of residual antimicrobial effect. Macro
25 and micro images were also taken to determine the levels of biofilm formation in the competitive product plates. The levels of biofilm formation in the Polymer S2/isopropanol treated plates and the competitive product treated plates were then compared. The sections of the polystyrene and glass surfaces treated with an isopropanol solution (0.25% w/v, 0.5% w/v, 1% w/v, 2% w/v) of the polymer S2
30 at a rate of 5 to 40 μ g/sq. in. showed virtually complete inhibition of *Klebsiella pneumonia* and *Pseudomonas fluorescens* biofilm formation whereas biofilm grew in the untreated sections of the polystyrene and glass surfaces. The competitive product treated plates had biofilm growing on both the treated and the untreated

sections of the plates 24 hours after treatment under the above test conditions. Thus, the Polymer S2/isopropanol solution provides residual inhibition against the growth of biofilm on the surfaces.

5 [0057] Therefore, there has been provided a hard surface antimicrobial cleaner which cleans and disinfects a hard surface and also provides for residual effectiveness versus bacteria. Further, there has been provided a hard surface antimicrobial cleaner which cleans and disinfects a hard surface and thereafter inhibits the formation of biofilm on the hard surface. Although the present invention has been described in detail with reference to certain embodiments, one
10 skilled in the art will appreciate that the present invention can be practiced by other than the described embodiments, which have been presented for purposes of illustration and not of limitation. Therefore, the scope of the appended claims should not be limited to the description of the embodiments contained herein.

INDUSTRIAL APPLICABILITY

15 [0058] The hard surface antimicrobial cleaner according to the invention may be used for cleaning and disinfecting a hard surface, and also provides for residual effectiveness versus bacteria on the hard surface and/or residual inhibition against the formation of biofilm on the hard surface.

CLAIMS

We claim:

1. A hard surface antimicrobial cleaner comprising:
 from about 0.01% to about 20% by weight of the total weight of the cleaner
 5 of a disinfectant;
 from about 0.01% to about 20% by weight of the total weight of the cleaner
 of a polysiloxane with at least one poly(oxyalkylene) side chain; and
 from about 0.5% to about 20% by weight of the total weight of the cleaner
 of a solvent,
 10 wherein the cleaner kills bacteria on a hard surface for at least 24 hours
 after being sprayed onto and wiped from the hard surface.

2. The cleaner of claim 1 wherein:
 the polysiloxane has the formula:
 15 $R_1R_2R_3SiO(R_4R_5SiO)_p(R_6QSiO)_qSiR_3R_2R_1$ in which R_1, R_2, R_4, R_5, R_6 are identical
 or different and are a C_1 - C_6 alkyl or phenyl, R_3 is identical or different and is C_1 - C_6
 alkyl, phenyl or Q, and Q is an ether polyoxyalkylene group of the formula –
 $R-O-(R'O)_nR''$ where R is a linear or branched C_3 - C_{15} alkyl group, $(R'O)_n$ is a
 poly(ethyleneoxy) and/or poly(propyleneoxy) group, n is a mean value ranging
 20 from 5 to 200, R'' is H or a C_1 - C_6 alkyl group, p is a mean value ranging from 10 to
 200, and q is 0 or a mean value ranging from 1 to 200, R_3 being Q when q is 0.

3. The cleaner of claim 1 wherein:
 the polysiloxane has the formula:
 25
- $$\begin{array}{ccccccc}
 & CH_3 & & CH_3 & & CH_3 & & CH_3 \\
 & | & & | & & | & & | \\
 H_3C - & Si & - & [Si-O]_6 & - & [Si-O]_{60} & - & Si - CH_3 \\
 & | & & | & & | & & | \\
 & CH_3 & & (CH_2)_3 & & CH_3 & & CH_3 \\
 & & & | & & & & \\
 & & & [O - CH_2 - CH_2]_{23} - [O - CH_2 - CH_2]_{22} - OH.
 \end{array}$$
- 30

4. The cleaner of claim 1 further comprising:
 from 0.1 to about 20% by weight of the total weight of the cleaner of a
 35 sequesterant.

5. The cleaner of claim 1 further comprising:
from 0.5 to about 20% by weight of the total weight of the cleaner of a
surfactant selected from nonionic surfactants, amphoteric surfactants, sarcosine
anionic surfactants, cationic surfactants and mixtures thereof.

6. The cleaner of claim 1 further comprising:
from 0.01 to about 5% by weight of the total weight of the cleaner of a
water soluble organosilane of the formula: $A_{3-x}B_xSiD$ wherein A is -OH or a
hydrolyzable group, B is an alkyl group of from 1 to 4 carbon atoms, x has a value
of 0, 1 or 2, and D is a hydrocarbon group of from 1 to 4 carbon atoms, phenyl, or
a nonionic or cationic, substituted-hydrocarbon group containing at least one
oxygen or nitrogen group or salts of such substituted-hydrocarbon groups.

7. A method for cleaning and disinfecting a hard surface and providing
residual effectiveness against bacteria to the hard surface, the method
comprising:
applying the cleaner of claim 1 to the hard surface.


8. A hard surface antimicrobial cleaner comprising:
from about 0.01% to about 20% by weight of the total weight of the cleaner
of a disinfectant;
from about 0.01% to about 20% by weight of the total weight of the cleaner
of a polysiloxane with at least one poly(oxyalkylene) side chain; and
from 0.1 to about 20% by weight of the total weight of the cleaner of a
sequesterant,
wherein the cleaner kills bacteria on a hard surface for at least 24 hours
after being sprayed onto and wiped from the hard surface.

9. The cleaner of claim 8 wherein:
the polysiloxane has the formula:
 $R_1R_2R_3SiO(R_4R_5SiO)_p(R_6QSiO)_qSiR_3R_2R_1$ in which R_1, R_2, R_4, R_5, R_6 are identical
or different and are a C_1 - C_6 alkyl or phenyl, R_3 is identical or different and is C_1 - C_6

alkyl, phenyl or Q, and Q is an ether polyoxyalkylene group of the formula –
 R-O-(R'O)_nR'' where R is a linear or branched C₃-C₁₅ alkyl group, (R'O)_n is a
 poly(ethyleneoxy) and/or poly(propyleneoxy) group, n is a mean value ranging
 from 5 to 200, R'' is H or a C₁-C₆ alkyl group, p is a mean value ranging from 10 to
 5 200, and q is 0 or a mean value ranging from 1 to 200, R₃ being Q when q is 0.

10. The cleaner of claim 8 wherein:

the polysiloxane has the formula:



11. The cleaner of claim 8 further comprising:

from 0.5 to about 20% by weight of the total weight of the cleaner of a
 surfactant selected from nonionic surfactants, amphoteric surfactants, sarcosine
 20 anionic surfactants, cationic surfactants and mixtures thereof.

12. The cleaner of claim 8 further comprising:

from 0.01 to about 5% by weight of the total weight of the cleaner of a
 water soluble organosilane of the formula: A_{3-x}B_xSiD wherein A is -OH or a
 25 hydrolyzable group, B is an alkyl group of from 1 to 4 carbon atoms, x has a value
 of 0, 1 or 2, and D is a hydrocarbon group of from 1 to 4 carbon atoms, phenyl, or
 a nonionic or cationic, substituted-hydrocarbon group containing at least one
 oxygen or nitrogen group or salts of such substituted-hydrocarbon groups.

30 13. A method for cleaning and disinfecting a hard surface and providing
 residual effectiveness against bacteria to the hard surface, the method
 comprising:

applying the cleaner of claim 8 to the hard surface.

35

14. A hard surface antimicrobial cleaner comprising:

from about 0.01% to about 20% by weight of the total weight of the cleaner of a disinfectant;

from about 0.01% to about 20% by weight of the total weight of the cleaner of a polysiloxane with at least one poly(oxyalkylene) side chain; and

5 from 0.5 to about 20% by weight of the total weight of the cleaner of a surfactant selected from nonionic surfactants, amphoteric surfactants, sarcosine anionic surfactants, cationic surfactants and mixtures thereof,

wherein the cleaner kills bacteria on a hard surface for at least 24 hours after being sprayed onto and wiped from the hard surface.

10

15. The cleaner of claim 14 wherein:


the polysiloxane has the formula:

15 $R_1R_2R_3SiO(R_4R_5SiO)_p(R_6QSiO)_qSiR_3R_2R_1$ in which R_1, R_2, R_4, R_5, R_6 are identical or different and are a C_1 - C_6 alkyl or phenyl, R_3 is identical or different and is C_1 - C_6 alkyl, phenyl or Q, and Q is an ether polyoxyalkylene group of the formula – $R-O-(R'O)_nR''$ where R is a linear or branched C_3 - C_{15} alkyl group, $(R'O)_n$ is a poly(ethyleneoxy) and/or poly(propyleneoxy) group, n is a mean value ranging from 5 to 200, R'' is H or a C_1 - C_6 alkyl group, p is a mean value ranging from 10 to 200, and q is 0 or a mean value ranging from 1 to 200, R_3 being Q when q is 0.

20

16. The cleaner of claim 14 wherein:

the polysiloxane has the formula:



30

17. The cleaner of claim 14 further comprising:

from 0.01 to about 5% by weight of the total weight of the cleaner of a water soluble organosilane of the formula: $A_{3-x}B_xSiD$ wherein A is -OH or a hydrolyzable group, B is an alkyl group of from 1 to 4 carbon atoms, x has a value of 0, 1 or 2, and D is a hydrocarbon group of from 1 to 4 carbon atoms, phenyl, or

35

a nonionic or cationic, substituted-hydrocarbon group containing at least one oxygen or nitrogen group or salts of such substituted-hydrocarbon groups.

18. A method for cleaning and disinfecting a hard surface and providing residual effectiveness against bacteria to the hard surface, the method comprising:

applying the cleaner of claim 14 to the hard surface.

19. A hard surface antimicrobial cleaner comprising:
from about 0.01% to about 20% by weight of the total weight of the cleaner of a disinfectant;

from about 0.01% to about 20% by weight of the total weight of the cleaner of a polysiloxane with at least one poly(oxyalkylene) side chain; and

a water soluble organosilane of the formula: $A_{3-x}B_xSiD$ wherein A is -OH or a hydrolyzable group, B is an alkyl group of from 1 to 4 carbon atoms, x has a value of 0, 1 or 2, and D is a hydrocarbon group of from 1 to 4 carbon atoms, phenyl, or a nonionic or cationic, substituted-hydrocarbon group containing at least one oxygen or nitrogen group or salts of such substituted-hydrocarbon groups,

wherein the cleaner kills bacteria on a hard surface for at least 24 hours after being sprayed onto and wiped from the hard surface.


20. The cleaner of claim 19 wherein:

the polysiloxane has the formula:

$R_1R_2R_3SiO(R_4R_5SiO)_p(R_6QSiO)_qSiR_3R_2R_1$ in which R_1, R_2, R_4, R_5, R_6 are identical or different and are a C_1 - C_6 alkyl or phenyl, R_3 is identical or different and is C_1 - C_6 alkyl, phenyl or Q, and Q is an ether polyoxyalkylene group of the formula – $R-O-(R'O)_nR''$ where R is a linear or branched C_3 - C_{15} alkyl group, $(R'O)_n$ is a poly(ethyleneoxy) and/or poly(propyleneoxy) group, n is a mean value ranging from 5 to 200, R'' is H or a C_1 - C_6 alkyl group, p is a mean value ranging from 10 to 200, and q is 0 or a mean value ranging from 1 to 200, R_3 being Q when q is 0.

21. The cleaner of claim 19 wherein:

the polysiloxane has the formula:



22. The cleaner of claim 19 wherein:
the water soluble organosilane is present in the cleaner in the range of from 0.01 to about 5% by weight of the total weight of the cleaner.

23. A method for cleaning and disinfecting a hard surface and providing residual effectiveness against bacteria to the hard surface, the method comprising:

applying the cleaner of claim 19 to the hard surface.

24. A hard surface antimicrobial cleaner comprising:
from about 50% to about 99.9% by weight of the total weight of the cleaner of an alkyl alcohol disinfectant; and
a polysiloxane with at least one poly(oxyalkylene) side chain,
wherein the cleaner inhibits biofilm formation on a hard surface for at least 24 hours after being sprayed onto the hard surface.

25. The cleaner of claim 24 wherein:
the disinfectant is present in the cleaner in the range of about 60% to about 80% by weight of the total weight of the cleaner.

26. The cleaner of claim 24 wherein:
the polysiloxane with at least one poly(oxyalkylene) side chain is present in the cleaner in the range of from about 0.01% to about 20% by weight of the total weight of the cleaner.


27. The cleaner of claim 24 wherein:

the polysiloxane has the formula:

$R_1R_2R_3SiO(R_4R_5SiO)_p(R_6QSiO)_qSiR_3R_2R_1$ in which R_1, R_2, R_4, R_5, R_6 are identical or different and are a C_1 - C_6 alkyl or phenyl, R_3 is identical or different and is C_1 - C_6 alkyl, phenyl or Q, and Q is an ether polyoxyalkylene group of the formula –
 5 $R-O-(R'O)_nR''$ where R is a linear or branched C_3 - C_{15} alkyl group, $(R'O)_n$ is a poly(ethyleneoxy) and/or poly(propyleneoxy) group, n is a mean value ranging from 5 to 200, R'' is H or a C_1 - C_6 alkyl group, p is a mean value ranging from 10 to 200, and q is 0 or a mean value ranging from 1 to 200, R_3 being Q when q is 0.

28. The cleaner of claim 24 wherein:

the polysiloxane has the formula:



29. The cleaner of claim 24 wherein:

the alkyl alcohol is ethanol, n-propanol or isopropanol.

30. The cleaner of claim 24 wherein:

the alkyl alcohol is isopropanol.

31. The cleaner of claim 24 wherein:

the alkyl alcohol is ethanol, n-propanol or isopropanol, and

the disinfectant is present in the cleaner in the range of about 60% to about 80% by weight of the total weight of the cleaner.

32. The cleaner of claim 24 wherein:

the alkyl alcohol is ethanol, n-propanol or isopropanol, and

the disinfectant is present in the cleaner in the range of about 60% to about 80% by weight of the total weight of the cleaner, and

the polysiloxane with at least one poly(oxyalkylene) side chain is present in the cleaner in the range of from about 0.01% to about 20% by weight of the total weight of the cleaner.

5 33. A method for cleaning and disinfecting a hard surface and providing residual inhibition against biofilm formation on the hard surface, the method comprising:

 applying the cleaner of claim 24 to the hard surface.

10 34. The method of claim 33 wherein:
 biofilm formation is inhibited for at least 24 hours.

INTERNATIONAL SEARCH REPORT

International Application No

PCT/US 02/07538

A. CLASSIFICATION OF SUBJECT MATTER

IPC 7 C11D11/00 C11D3/00 C11D3/37 C11D3/20 C11D7/26
C11D3/33

According to International Patent Classification (IPC) or to both national classification and IPC

B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

IPC 7 C11D

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practical, search terms used)

EPO-Internal, WPI Data, PAJ

C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	WO 99 18784 A (CARR ET AL) 22 April 1999 (1999-04-22) cited in the application page 2, line 7-12	1-24
A	page 4, line 1-5; claim 1 page 7, line 8 example 3	24-34
X	FR 2 771 416 A (RHONE POULENC CHIMIE) 28 May 1999 (1999-05-28) page 8, line 27 - line 29 example B claim 1	1-3,5,7, 14-16,18

	---/---	



Further documents are listed in the continuation of box C.



Patent family members are listed in annex.

* Special categories of cited documents:

- *A* document defining the general state of the art which is not considered to be of particular relevance
- *E* earlier document but published on or after the international filing date
- *L* document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)
- *O* document referring to an oral disclosure, use, exhibition or other means
- *P* document published prior to the international filing date but later than the priority date claimed

- *T* later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention
- *X* document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone
- *Y* document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art.
- *G* document member of the same patent family

Date of the actual completion of the international search

22 July 2002

Date of mailing of the international search report

31/07/2002

Name and mailing address of the ISA

European Patent Office, P.B. 5818 Patentlaan 2
NL - 2280 HV Rijswijk
Tel. (+31-70) 340-2040, Tx. 31 651 epo nl,
Fax: (+31-70) 340-3016

Authorized officer

Rasmusson, R

INTERNATIONAL SEARCH REPORT

International Application No
PCT/US 02/07538

C.(Continuation) DOCUMENTS CONSIDERED TO BE RELEVANT		
Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
A	US 6 087 319 A (NORMAN CLARE L) 11 July 2000 (2000-07-11) cited in the application column 1, line 9 - line 19 column 2, line 9 -column 3, line 25 claims 1,6 ---	1-34
X	US 3 671 630 A (CARROLL ET AL) 20 June 1972 (1972-06-20) column 2, line 54 - line 56 example XXVII -----	24-34

INTERNATIONAL SEARCH REPORT

 International Application No
 PCT/US 02/07538

Patent document cited in search report		Publication date		Patent family member(s)	Publication date
WO 9918784	A	22-04-1999	FR	2769469 A1	16-04-1999
			AU	9545298 A	03-05-1999
			CA	2305496 A1	22-04-1999
			EP	1022941 A1	02-08-2000
			WO	9918784 A1	22-04-1999
FR 2771416	A	28-05-1999	FR	2771416 A1	28-05-1999
			AU	1162499 A	15-06-1999
			WO	9927031 A1	03-06-1999
US 6087319	A	11-07-2000	AU	718739 B2	20-04-2000
			AU	2055497 A	22-09-1997
			CA	2248219 A1	12-09-1997
			EP	0885280 A1	23-12-1998
			JP	2000506209 T	23-05-2000
			NZ	331796 A	28-10-1999
			WO	9732957 A1	12-09-1997
			ZA	9701858 A	24-10-1997
US 3671630	A	20-06-1972	FI	51367 A	